

$K_{\ell 3}$ and π_{e3} transition form factorsYu. Kalinovsky,^a K. L. Mitchell^b and C. D. Roberts^c^a *Bogoliubov Laboratory of Theoretical Physics,
Joint Institute for Nuclear Research, 141980 Dubna, Russia*^b *TRIUMF, 4004 Wesbrook Mall,
Vancouver, British Columbia, Canada V6T 2A3*^c *Physics Division, Bldg. 203, Argonne National Laboratory,
Argonne IL 60439-4843, USA*

Abstract

$K_{\ell 3}$ and π_{e3} transition form factors are calculated as an application of Dyson-Schwinger equations. The role of nonanalytic contributions to the quark-W-boson vertex is elucidated. A one-parameter model for this vertex provides a uniformly good description of these transitions, including the value of the scalar form factor of the kaon at the Callan-Treiman point. The $K_{\ell 3}$ form factors, $f_{\pm}^K$, are approximately linear on $t \in [m_e^2, m_{\mu}^2]$ and have approximately the same slope. $f_{-}^K(0)$ is a measure of the Euclidean constituent-quark mass ratio: M_s^E/M_u^E . In the isospin symmetric limit: $-f_{+}^{\pi}(0) = F_{\pi}(t)$, the electromagnetic pion form factor, and $f_{-}^{\pi}(t) \equiv 0$.

Keywords: Electroweak interactions; Semileptonic decays, $f_{K_{\ell 3}}(t)$, $f_{\pi_{e3}}(t)$; Dyson-Schwinger equations; Confinement; Nonperturbative QCD; Quark models.

1. Introduction.

The semileptonic transitions $K^+ \rightarrow \pi^0 \ell \nu_{\ell}$ [$K_{\ell 3}^+$], $K^0 \rightarrow \pi^- \ell \nu_{\ell}$ [$K_{\ell 3}^0$] and $\pi^+ \rightarrow \pi^0 e \nu_e$ [π_{e3}] proceed via the flavour-changing, vector piece of the $V - A$ electroweak interaction, in particular j_{μ}^{su} and j_{μ}^{du} . The axial-vector component does not contribute because, in every case, the two mesons involved have the same parity. Neither j_{μ}^{su} nor j_{μ}^{du} is conserved; in each case the symmetry breaking term is a measure of the current-quark mass difference, $m_s - m_u$ or $m_d - m_u$, and its enhancement due to nonperturbative effects. Therefore these processes can be employed to probe $SU_f(3)$ flavour symmetry violation.

The current status of experimental analyses of $K_{\ell 3}^+$ and $K_{\ell 3}^0$ transitions is summarised in Ref. [1]. It is not completely satisfactory, with the value of two of the four observables showing a surprising sensitivity to the initial state (see

Table 2). Contemporary theoretical analyses fall into two classes: those employing quark-gluon degrees of freedom; e.g., Refs. [2,3], and those using meson degrees of freedom; e.g., Refs. [4,5]. While there is agreement between these approaches for some observables, in Sect. 3 we identify *qualitatively important* quantitative disagreements whose resolution requires an appreciation of the origin and role of nonanalytic contributions to the quark-W-boson vertex.

Our analysis of the transition amplitudes, and their form factors, is a phenomenological application of Dyson-Schwinger equations [DSEs]. It is an extension of the calculation of the electromagnetic form factors: $F_{K^\pm}(t)$, $F_{K^0}(t)$ [6] and $F_\pi(t)$ [7]. As therein, primary elements of this calculation are the dressed-quark propagator (2-point Schwinger function) for the u -, d - and s -quarks and the Bethe-Salpeter amplitudes for the π - and K -mesons, the behaviour of which follows from extensive nonperturbative, model-DSE studies [8].

A significant new feature of this study is the dressed-vertex (3-point Schwinger function) describing the quark- W -boson coupling. A qualitative understanding of this is crucial in resolving the discrepancies between quark- and meson-based analyses of the transition form factors; and is an important precursor to the study of other weak-interaction processes such as those involving baryons.

2. Transition Form Factors.

The matrix elements for the $K_{\ell 3}$ and $\pi_{e 3}$ transitions are

$$J_\mu^{K^+}(K, Q) \equiv \langle \pi^0(p) | \bar{u} \gamma_\mu s | K^+(k) \rangle \equiv \frac{1}{\sqrt{2}} \left(f_+^{K^+}(t) K_\mu + f_-^{K^+}(t) Q_\mu \right) \quad (1)$$

$$J_\mu^{K^0}(K, Q) \equiv \langle \pi^-(p) | \bar{u} \gamma_\mu s | K^0(k) \rangle \equiv f_+^{K^0}(t) K_\mu + f_-^{K^0}(t) Q_\mu \quad (2)$$

$$J_\mu^\pi(K, Q) \equiv \langle \pi^0(p) | \bar{u} \gamma_\mu d | \pi^+(k) \rangle \equiv \frac{1}{\sqrt{2}} \left(f_+^\pi(t) K_\mu + f_-^\pi(t) Q_\mu \right), \quad (3)$$

where $K = k + p$, $Q = k - p$ and the squared-momentum transfer $t = -Q^2$.¹

In the isospin-symmetric case, $m_u = m_d$, j_μ^{du} is conserved and: 1) $f_\pm^{K^0} \equiv f_\pm^{K^+}$; 2) $f_+^\pi(t) = -F_\pi(t)$; and 3) $f_-^\pi \equiv 0$, while in the case of $SU_f(3)$ symmetry, j_μ^{su} is also conserved, $f_+^{K^0, K^+, \pi}(t) \equiv -F_\pi(t)$ and $f_-^{K^0, K^+, \pi} \equiv 0$. Away from the symmetry limits, the Ademollo-Gatto theorem [10] entails $f_+^2(0) \approx 1$; i.e., in the vector form factor, $SU_f(3)$ symmetry breaking effects are suppressed at $t = 0$. However, one expects $f_-(0)$ to be sensitive to the nonperturbative enhancement of the $SU_f(3)$ symmetry breaking current-quark mass differences, since $f_-(0)$ depends on the $s:u$ ratio of constituent-quark masses in Ref. [11].

¹ We employ a Euclidean space formulation with $\{\gamma_\mu, \gamma_\nu\} = 2\delta_{\mu\nu}$, $\gamma_\mu^\dagger = \gamma_\mu$ and $a \cdot b = \sum_{i=1}^4 a_i b_i$. A timelike vector, Q_μ , has $Q^2 < 0$.

The scalar form factor

$$f_0^K(t) \equiv f_+^K(t) + \frac{t}{m_K^2 - m_\pi^2} f_-^K(t) \quad (4)$$

measures the divergence of the currents in Eqs. (1) and (2), which makes it a particularly useful observable. Current algebra predicts the value of f_0^K at the Callan-Treiman point, $t = m_K^2 - m_\pi^2 \equiv \Delta$: $f_0^K(\Delta) = -f_K/f_\pi$, the ratio of the weak-decay constants [12]. The Callan-Treiman point is not accessible experimentally, however, the robust nature of the derivation of this result makes it a useful tool in constraining and improving a given theoretical framework.

In impulse approximation

$$J_\mu^{K^+}(K, Q) = \sqrt{2}N_c \int \frac{d^4\ell}{(2\pi)^4} \text{tr}_D \left[\bar{\Gamma}_{\pi^0} \left(\ell + \frac{1}{4}[K + Q]; -p \right) \times \right. \\ \left. S_u(\ell + \frac{1}{2}K) \Gamma_{K^+} \left(\ell + \frac{1}{4}[K - Q]; k \right) S_s(\ell - \frac{1}{2}Q) iV_\mu^{su}(\ell; -Q) S_u(\ell + \frac{1}{2}Q) \right], \quad (5)$$

with obvious modifications for $K_{\ell 3}^0$ and π_{e3} . In Eq. (5): S_f is the dressed-propagator for quark flavour f ; $\Gamma_M(q; P)$ is the Bethe-Salpeter amplitude for a meson M , with q the relative quark-antiquark momentum and P the total momentum; and $V_\mu^{su}(q; P)$ is the dressed s - u - W -vertex. Herein we work in the isospin symmetric limit; i.e., we do not distinguish between u - and d -quarks, and hence $S_u \equiv S_d$.

Extensive studies of the DSE for the dressed-quark propagator [8] lead to the following algebraic approximating form, used successfully in Refs. [6,9]: $S_f^f(p) = -i\gamma \cdot p \sigma_V^f(p) + \sigma_S^f(p)$,

$$\bar{\sigma}_S^f(x) = C^f(\bar{m}_f) e^{-2x} + \frac{\bar{m}_f}{x + \bar{m}_f^2} \left(1 - e^{-2(x + \bar{m}_f^2)} \right) \quad (6)$$

$$+ \frac{1 - e^{-b_1^f x}}{b_1^f x} \frac{1 - e^{-b_3^f x}}{b_3^f x} \left(b_0^f + b_2^f \frac{1 - e^{-\Lambda x}}{\Lambda x} \right) \\ \bar{\sigma}_V^f(x) = \frac{2(x + \bar{m}_f^2) - 1 + e^{-2(x + \bar{m}_f^2)}}{2(x + \bar{m}_f^2)^2} - \bar{m}_f C^f(\bar{m}_f) e^{-2x}, \quad (7)$$

where $x = p^2/(2D)$ and: $\bar{\sigma}_V^f(x) = 2D \sigma_V^f(p^2)$; $\bar{\sigma}_S^f(x) = \sqrt{2D} \sigma_S^f(p^2)$; and $\bar{m}_f = m_f/\sqrt{2D}$, with D a mass scale. We write the inverse of the propagator as $S_f^{-1}(p) = i\gamma \cdot p A^f(p^2) + B^f(p^2)$.

The Bethe-Salpeter equation [BSE] for the pseudoscalar mesons has also been studied extensively [8]. The realisation of Goldstone's theorem entails that, in the chiral limit, $m_f = 0$, the pseudoscalar Bethe-Salpeter amplitude is

completely determined by the dressed-quark propagator [13]. As a consequence one finds that

$$\Gamma_\pi(p; P^2 = -m_\pi^2) \approx i\gamma_5 \frac{1}{f_\pi} B_{m_u=0}^u(p^2), \quad (8)$$

$$\Gamma_K(p; P^2 = -m_K^2) \approx i\gamma_5 \frac{1}{f_K} B_{m_s=0}^s(p^2), \quad (9)$$

with $f_\pi \approx 92$ MeV and $f_K \approx 113$ MeV the calculated pion and kaon normalisation constants, respectively, are good pointwise approximations. These results follow from BSE studies and have proven phenomenologically efficacious [6,9].

The parameters $C^f(\bar{m}_f)$, $\bar{m}_f$, $b_{1...3}^f$ in Eqs. (6), (7) are determined in a χ^2 -fit to a range of hadronic observables. This is described in Ref. [6] and leads to the values in Table 1. $C^f(\bar{m}_f)$ is a function of $\bar{m}_f$; it is only non-zero when evaluating the Bethe-Salpeter amplitudes using Eqs. (8), (9), which allows the algebraic approximation to represent those differences between $B_{m=0}(0)$ and $B_{m \neq 0}(0)$ observed in numerical studies of the quark DSE. In the fit, the difference between the u - and d -quarks was neglected and only that minimal difference between u - and s -quarks allowed that was necessary to ensure: $\langle \bar{s}s \rangle < \langle \bar{u}u \rangle$; and $m_s/m_u \gg 1$. ($\Lambda = 10^{-4}$ is included in Eqs. (6), (7) only for the purpose of separating the small- and intermediate- p^2 behaviour of the algebraic form, characterised by b_0 and b_2 ; a separation in magnitude observed in numerical studies.)

Table 1

The values of $b_{1,3}^s$ are underlined to indicate that the constraints $b_{1,3}^s = b_{1,3}^u$ were imposed in the fitting. The scale parameter $D = 0.160$ GeV².

	$C^f(\bar{m}_f = 0)$	$\bar{m}_f$	b_0^f	b_1^f	b_2^f	b_3^f
u	0.121	0.00897	0.131	2.90	0.603	0.185
s	1.69	0.224	0.105	<u>2.90</u>	0.740	<u>0.185</u>

The algebraic approximation represented by Eqs. (6)-(9) provides a good description of pion and kaon observables [6] and has recently been employed successfully in the prediction of a wide range of vector-meson electroproduction observables [9]; a process far outside the domain on which it was constrained.

2.1 Dressed Quark W -Boson Vertex. $V_\mu^{su}(p; P)$ in Eq. (5) is the new element in this study. It satisfies a DSE, from which one can derive the following Ward-Takahashi identity

$$Q_\mu iV_\mu^{su}(p; Q) = S_s^{-1}(p_+) - S_u^{-1}(p_-) - (m_s - m_u) \Gamma_I^{su}(p; Q) \quad (10)$$

where $p_\pm \equiv p \pm Q/2$ and $\Gamma_I^{su}(p; Q)$ is the flavour-dependent scalar vertex (neglecting interactions, $\Gamma_I^{su}(p; Q) = I_D$). This entails that the flavour-changing

vector current is not conserved and that the symmetry breaking term measures the current-quark mass difference. Nonperturbative contributions to $\Gamma_I^{su}(p; Q)$ enhance this effect.

Herein we use what is known from extensive studies of the flavour-dependent, electromagnetic quark-photon vertex, $\Gamma_\mu^f(p; Q)$, to construct an Ansatz for $V_\mu^{su}(p; P)$. In studies of pion and kaon electromagnetic form factors [6,7] and vector meson electroproduction [9] the Ball-Chiu Ansatz [14]

$$i\Gamma_\mu^f(p; Q) = i\gamma_\mu f_1^f(p; Q) + i\gamma \cdot p p_\mu f_2^f(p; Q) + p_\mu f_3^f(p; Q), \quad (11)$$

where

$$\begin{aligned} f_1^f(p; Q) &= [A^f(p_+) + A^f(p_-)]/2, \\ f_2^f(p; Q) &= [A^f(p_+) - A^f(p_-)]/p \cdot Q, \\ f_3^f(p; Q) &= [B^f(p_+) - B^f(p_-)]/p \cdot Q \end{aligned} \quad (12)$$

has proven successful. As described in Ref. [7], this Ansatz is successful because it is the minimal solution of the Abelian Ward-Identity that is completely determined by the dressed-quark propagator and: A) has a well defined limit as $Q^2 \rightarrow 0$; B) transforms correctly under C , P , T and Lorentz transformations; and C) reduces to the bare vertex in the manner prescribed by perturbation theory. It therefore satisfies four of the six physical constraints proposed in Ref. [15] and explored in detail in Ref. [16]. Consequently, a first, minimal Ansatz is

$$V_\mu^{su}(p; Q) = \frac{1}{2} \left(\Gamma_\mu^s(p; Q) + \Gamma_\mu^u(p; Q) \right), \quad (13)$$

which satisfies Eq. (10), with a particular form of the symmetry breaking term, and also A)-C) above.

A property that our study, when using Eq. (13), shares with other quark-based studies; for example, Refs. [2,3,11], is that the s - u - W -vertex is analytic on the real- t axis. An Ansatz with this property excludes resonance contributions, such as K - π loops, which although unimportant for $t < 0$, may provide significant contributions for $t \in [m_e^2, m_\mu^2]$. We now address this issue.

3. Results and Discussion.

Equations (5)-(9) and (13) yield the results in the first column of Table 2. They are in quantitative agreement with the results of Refs. [2,3,11] and, with some *qualitatively important* exceptions that we discuss below, with Refs. [4,5].

Before addressing these exceptions we discuss other points of interest. The result for $f_+(0)$ is consistent with the Ademollo-Gatto theorem. Further, the

calculated value of $f_-(t_m)$ can be compared with Ref. [11]. Using our dressed-quark propagators we define a Euclidean constituent-quark mass, M_f^E , [17] which provides a single, indicative and quantitative measure of the importance of nonperturbative dressing of the quarks in the infrared: $(M_f^E)^2$ is the solution of $p^2[A^f(p^2)]^2 - [B^f(p^2)]^2 = 0$. In the present case we find $M_{u,d}^E = 330$ MeV, $M_s^E = 490$ MeV. From Fig. 1 of Ref. [11] these values lead to $f_-(t_m) \sim 0.28$, consistent with our result. (There is a relative “−” sign between our definitions of $f_\pm$ and those of Ref. [11].) This emphasises that $f_-(0)$ is a probe of the nonperturbative enhancement of $SU_f(3)$ breaking.

In the experimental analysis of $K_{\ell 3}$ decays it is often assumed that, on $t \in [m_e^2, m_\mu^2]$, $f_+(t)$ is linear; i.e., $f_+(t) = f_+(0)[1 + \lambda_+ t/m_\pi^2]$, and $f_-(t) = f_-(0)[1 + \lambda_- t/m_\pi^2]$, with $\lambda_- = 0$; i.e., $f_-(t) = \text{constant}$. Using the definition of λ in Table 2 we find that the difference $\lambda_+^e - \lambda_+^\mu$ is small and hence that a linear approximation to f_+ is reasonable. However, in our study we find $\lambda_- \sim \lambda_+$, consistent with Refs.[2,3] where $\lambda_+ = \lambda_-$, but inconsistent with the assumption that $f_-(t) = \text{constant}$.

In addressing the exceptions, we note from Table 2 that, in contrast to Ref. [5], neither our calculation using Eq. (13) nor Refs.[2,3] reproduce the anticipated value of the scalar form factor at the Callan-Treiman point, $f_0(\Delta) = -f_K/f_\pi$. Correlated with this are the results for λ_0 : in comparison with the experimental analysis of $K_{\mu 3}^0$ decays, which have the smallest error, λ_0 is of the wrong-sign or, if of the correct sign, it is an order-of-magnitude too small. These results are systematic effects due to the magnitude of $f_\pm(0)$ and the fact that $\lambda_+ \simeq \lambda_-$. To establish this we employed many variants of Eq. (13), the simplest of which replaces $[\Gamma_\mu^s + \Gamma_\mu^u]/2$ by $\alpha\Gamma_\mu^s + (1 - \alpha)\Gamma_\mu^u$, $0 < \alpha < 1$. The various Ansätze could introduce quantitative changes of $\sim 5\%$ in $f_0(\Delta)$ and $\sim 50\%$ in λ_0 , however, such changes are patently inadequate.

3.1 Nonanalytic Contributions to the Vertex. This discrepancy is a defect of the vertex Ansatz, Eq. (13); a defect that is implicit in Refs. [2,3] and similar studies. That this is the case is signalled by the small value of $r_{\pi K}$ in the first column of Table 2. Nonanalytic, resonance contributions to the quark-photon vertex or, analogously, π - π rescattering, provide $\lesssim 10\%$ of r_π [18], which measures $F'_\pi(t=0)$. Such nonanalytic contributions rapidly becomes insignificant as $(-t)$ increases because resonances do not contribute to the vertex at space-like momentum transfer [19]. However, their importance increases in the time-like region and, in calculating $f_0(t = \Delta \approx 0.56(m_K + m_\pi)^2)$, one is nearing the K - π production threshold where such contributions dominate.

In a careful study of the DSE for the s - u - W -vertex, these contributions are manifest in the solution at $t > 0$. They should therefore be included in constructing a realistic Ansatz for V_μ^{su} and hence, following Ref. [18], we consider

Table 2

$t_m = (m_K - m_\pi)^2$ is the largest, physically accessible value of the squared momentum-transfer, t ; $\xi(t) \equiv f_-(t)/f_+(t)$; $\lambda_\alpha^\ell \equiv m_{\pi^+}^2 f'_\alpha(m_\ell^2)/f_\alpha(0)$, $\alpha \in \{+, -, 0\}$; and $r_{\pi K}^2 \equiv 6f'_+(0)/f_+(0)$. Experimental results are taken from Ref. [1]. $m_e^2/m_\mu^2 \ll 1$ means that λ_0^e is not accessible experimentally. References to theoretical comparisons are labelled: a = Ref. [2], b = Ref. [3], c = Ref. [4], d = Ref. [5]; where necessary, results have been calculated using the information provided.

	Eq. (13)	Eq. (14)	Comparison	Experiment: $K_{\mu 3}^+$ $K_{\mu 3}^0$
$-f_+(t_m)$	1.11	1.24	1.04^a	
$f_-(t_m)$	0.27	0.30	0.29^a	
$-f_+(0)$	0.98	0.98	$0.93^a, 0.98^c$	
$f_-(0)$	0.24	0.24	$0.26^a, 0.15^d$	
$-\xi(0)$	0.25	0.25	$0.28^a, 0.28^b, 0.15^d$	0.35 ± 0.15 0.11 ± 0.09
λ_+^e	0.018	0.030	$0.019^a, 0.028^b, 0.030^d$	0.0286 ± 0.0022 0.0300 ± 0.0016
λ_+^μ	0.018	0.031	$0.019^a, 0.028^b, 0.030^d$	0.033 ± 0.008 0.034 ± 0.005
λ_-^e	0.012	0.024	$0.019^a, 0.028^b,$	
λ_-^μ	0.012	0.025	$0.019^a, 0.028^b,$	
$-f_0(\Delta)$	0.95	1.22	$0.88^a, 1.22^d$	
λ_0^e	-0.0024	0.0082	$-0.005^a, 0.0026^b, 0.017^d$	
λ_0^μ	-0.0024	0.0089	$-0.005^a, 0.0026^b, 0.017^d$	0.004 ± 0.007 0.025 ± 0.006
$r_{\pi K} \text{ (fm)}$	0.47	0.60	$0.48^a, 0.60^d$	

a heuristic minimal modification of Eq. (13), redefining f_1 as follows:

$$f_1(k; Q) \rightarrow f_1(k; Q) \mathcal{M}(t), \quad \mathcal{M}(t) \equiv \left[1 + c_l e^{t/t_p} \mathcal{L}(t) \right], \quad (14)$$

with $t_p = (m_K + m_\pi)^2$, and where

$$\mathcal{L}(t) \equiv 2 + \ln \left[\frac{m_\pi^2}{m_K^2} \right] \left(\frac{\Delta}{t} - \frac{\Sigma}{\Delta} \right) - \frac{\nu(t)}{t} \ln \left[\frac{(t + \nu(t))^2 - \Delta^2}{(t - \nu(t))^2 - \Delta^2} \right], \quad (15)$$

with $\Sigma = m_K^2 + m_\pi^2$ and $\nu(t)^2 = (t - t_p)(t - t_m)$, expresses the essence of the nonanalytic structure of the K - π loop [5]. In Eq. (14), c_l parametrises the

relative weight of the analytic and nonanalytic terms at $t = \Delta$. Its value is determined by the requirement that this Ansatz yield $f_0(\Delta) = -f_K/f_\pi$. This Ansatz, which herein is only sampled on the domain $t \in (-\infty, \Delta]$, is only a useful heuristic tool if the required value of c_l is small, for then it satisfies our physical requirements: it leaves the vertex unmodified for spacelike- t ; and it has a logarithmic branch point at the K - π production threshold.

The results obtained using Eq. (14), with $c_l = 0.17$, are presented in the second column of Table 2. With this value of c_l , $0.94 < \mathcal{M}(t < 0) \leq 1$; i.e., for $t < 0$, the modification of the vertex is small. The value of the slope parameters, $\lambda_{+,0}$, are in agreement with experiment and Ref. [5], as is $r_{\pi K}$. As expected, $f_\pm(0)$ and $\xi(0)$ are unchanged. This outcome is qualitatively insensitive to the exact form of $\mathcal{L}$; for example, using $\mathcal{L}(t) = -\{1 + (t_p/t) \ln[1 - t/t_p]\}$ with $c_l = 0.335$, leads only to a $\sim 3\%$ reduction in $f_\pm(t_m)$ and a $\sim 27\%$ reduction in λ_0 .




Fig. 1. $f_+^K(Q^2)$: solid line, Eq. (14); dashed line, Eq. (13). f_+^K is approximately linear for $-t_m < Q^2 < 0$.

In Figs. 1 and 2, to further illustrate the effect of nonanalytic contributions, we plot the $K_{\ell 3}$ transition form factors. $K_{\ell 3}^+$ is not distinguished from $K_{\ell 3}^0$ because hitherto we have made no attempt to fine-tune isospin breaking effects.

In Fig. 3 we compare $f_+^K(Q^2)$ with $f_+^\pi(Q^2)$. Since we work in the isospin symmetric limit, $f_+^\pi(Q^2) = -F_\pi(Q^2)$. The results are qualitatively similar to those obtained in Ref. [3], although, as in Ref. [6], our results for $f_+^{K\pi}(Q^2)$ are uniformly smaller in magnitude.

4. Summary and Conclusions.

We have analysed the $K_{\ell 3}$ and π_{e3} transition form factors. An important new element of this study is the development of an understanding of the form of the dressed s - u - W -vertex, $V_\mu^{su}(k; Q)$, on the domain $-(m_K + m_\pi)^2 < Q^2 < \infty$. A simple heuristic Ansatz for V_μ^{su} , with one adjustable parameter, yields a




Fig. 2. $f_-^K(Q^2)$: solid line, Eq. (14); dashed line, Eq. (13). f_-^K is approximately linear for $-t_m < Q^2 < 0$ and it varies almost as rapidly as f_+^K .




Fig. 3. $-Q^2 f_+^K(Q^2)$: solid line; $-Q^2 f_+^\pi(Q^2) = Q^2 F_\pi(Q^2)$: dot-dashed line. The peak in $Q^2 f_+^K(Q^2)$, most pronounced for the kaon, is a characteristic signal of quark-antiquark recombination into the meson final state in exclusive processes.

uniformly good description of $K_{\ell 3}$ observables, as illustrated in column two of Table 2.

We identified a systematic discrepancy between the studies of Refs. [2,3] and Refs. [4,5] arising because, in Refs. [2,3], nonanalytic contributions to V_μ^{su} in the timelike region are overlooked. These contributions are important on the domain $0 \lesssim t \lesssim (m_K^2 - m_\pi^2)$ and allow for a correct description of the scalar form factor, f_0^K , in this region; for example, without these contributions it is not possible to reproduce the current-algebra prediction for $f_0^K(t)$ at the

Callan-Treiman point: $t = m_K^2 - m_\pi^2$. The difference between the results in columns one and two of Table 2 is a measure of the relative importance of these contributions to a given quantity.

Our results are consistent with the Ademollo-Gatto theorem, $f_+^2(0) \approx 1$. They also support the expectation [11] that $f_-^K(0)$ is a quantitative measure of the nonperturbative enhancement of the current-quark mass difference $m_s - m_u$; i.e., it is a measure of the Euclidean constituent-quark mass-ratio: M_s^E/M_u^E .

Figures 1 and 2 illustrate that it is a good approximation to consider $f_+(t)$ to be linear functions on the domain $0 < t < (m_K - m_\pi)^2$. In common with other studies that employ quark-gluon degrees-of-freedom, we find $\lambda_- \simeq \lambda_+$. This suggests that the assumption, employed in experimental analyses, that $f_-(t)$ is a constant on this domain, should be reconsidered.

A strength of the DSE framework is that it allows the study of the t -dependence of the transition form factors on the entire domain $-\infty < t < (m_K - m_\pi)^2$. In the isospin symmetric limit, $m_d = m_u$, $f_+^\pi(t) = -F_\pi(t)$, the electromagnetic pion form factor and $f_-^\pi(t) \equiv 0$. One also has $f_+^{K^+}(t) = f_+^{K^0}(t)$, however, $f_+^K(t) \neq -F_K(t)$, the electromagnetic kaon form factor. There are qualitative similarities; for example, $-tf_+^K(t)$ exhibits the peak characteristic of quark-antiquark recombination into the meson final state in this exclusive process, however, $|f_+^K(t)|$ falls-off less rapidly than $|F_K(t)|$ at large- $(-t)$ and this is a measure of the difference between the pion and kaon Bethe-Salpeter amplitudes; i.e., of nonperturbative effects in QCD.

Acknowledgments. K.L.M. and Yu.K. are grateful for the hospitality of the Physics Division at ANL during visits in which this study was conceived and conducted. The work of K.L.M was supported in part by the Natural Sciences and Engineering Research Council of Canada; that of C.D.R. by the US Department of Energy, Nuclear Physics Division, under contract number W-31-109-ENG-38. The calculations described herein were carried out using the resources of the National Energy Research Supercomputer Center.

References

- [1] Particle Data Group, Phys. Rev. D **50** (1994) 1173.
- [2] D. Scora and N. Isgur, Phys. Rev. D **52** (1992) 2783.
- [3] A. Afanasev and W. W. Buck, *Unified description of kaon electroweak form factors*, archive: hep-ph/9606296.
- [4] H. Leutwyler and M. Roos, Z. Phys. C **25** (1984) 91.
- [5] J. Gasser and H. Leutwyler, Nucl. Phys. B **250** (1985) 517.

- [6] C. J. Burden, C. D. Roberts and M. J. Thomson, Phys. Lett. B **371** (1996) 163.
- [7] C. D. Roberts, Nucl. Phys. A **605** (1996) 475.
- [8] C. D. Roberts and A. G. Williams, Prog. Part. Nucl. Phys. **33** (1994) 477.
- [9] M. A. Pichowsky and T. S. -H. Lee, Phys. Lett. B **379**, 1 (1996); *Exclusive, diffractive processes and the quark substructure of mesons*, preprint ANL-PHY-8529-TH-96.
- [10] N. Ademollo and R. Gatto, Phys. Rev. Lett. **13** (1964) 264.
- [11] N. Isgur, Phys. Rev. D **12** (1975) 3666.
- [12] C. G. Callan and S. B. Treiman, Phys. Rev. Lett. **16** (1966) 153.
- [13] A. Bender, C. D. Roberts and L. v. Smekal, Phys. Lett. B **380** (1996) 7; C. D. Roberts, *Confinement, Diquarks and Goldstone's theorem*, to appear in the proceedings of *Quark Confinement and the Hadron Spectrum, II* (World Scientific, Singapore), archive: nucl-th/9609039.
- [14] J. S. Ball and T.-W. Chiu, Phys. Rev. D **22** (1980) 2542.
- [15] C. J. Burden and C. D. Roberts, Phys. Rev. D **47** (1993) 5581.
- [16] A. Bashir and M. R. Pennington, Phys. Rev. D **50** (1994) 7679.
- [17] M. R. Pennington and D. Walsh, Phys. Lett. B **253** (1991) 246.
- [18] R. Alkofer, A. Bender and C. D. Roberts, Int J. Mod. Phys. A **10** (1995) 3319.
- [19] K. L. Mitchell and P.C. Tandy, *Pion loop contribution to ρ - ω mixing and mass splitting*, archive: nucl-th/9607025.